arXiv:1107.5879v2 [hep-ph] 8 Aug 2011

A Higgs Quadruplet for Type III Seesaw and Implications for $\mu \to e\gamma$ and $\mu - e$ Conversion

Bo Ren[1], Koji Tsumura[2], Xiao-Gang He[1,2],
[1]*INPAC, Department of Physics, Shanghai Jiao Tong University, Shanghai, China*
[2]*Department of Physics and Center for Theoretical Sciences,*
National Taiwan University, Taipei, Taiwan

(Dated: August 9, 2011)

In Type III seesaw model the heavy neutrinos are contained in leptonic triplet representations. The Yukawa couplings of the triplet fermion and the left-handed neutrinos with the doublet Higgs field produce the Dirac mass terms. Together with the Majorana masses for the leptonic triplets, the light neutrinos obtain non-zero seesaw masses. We point out that it is also possible to have a quadruplet Higgs field to produce the Dirac mass terms to facilitate the seesaw mechanism. The vacuum expectation value of the quadruplet Higgs is constrained to be small by electroweak precision data. Therefore the Yukawa couplings of a quadruplet can be much larger than those for a doublet. We also find that unlike the usual Type III seesaw model where at least two copies of leptonic triplets are needed, with both doublet and quadruplet Higgs representations, just one leptonic triplet is possible to have a phenomenologically acceptable model because light neutrino masses can receive sizable contributions at both tree and one loop levels. Large Yukawa couplings of the quadruplet can induce observable effects for lepton flavor violating processes $\mu \to e\gamma$ and $\mu - e$ conversion. Implications of the recent $\mu \to e\gamma$ limit from MEG and also limit on $\mu - e$ conversion on Au are also given. Some interesting collider signatures for the doubly charged Higgs boson in the quadruplet are discussed.

I. INTRODUCTION

The type III seesaw contains leptonic triplets Σ_R under the standard model (SM) gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ as $(1,3,0)$, $\Sigma_R = (\Sigma_R^+, \Sigma_R^0, \Sigma_R^-)$[1]. In tensor notation, the triplet can be written as $\Sigma_R = (\Sigma_{ij})$ symmetric in i and j, where i and j take the values 1 and 2. $\Sigma_{R11} = \Sigma_R^+$, $\Sigma_{R12} = i\Sigma_R^0/\sqrt{2}$ and $\Sigma_{R22} = \Sigma_R^-$. The Yukawa couplings related to neutrino and charged lepton masses come from the following terms

$$L = -\bar{L}_L Y_e E_R \Phi - \bar{L}_L Y_\nu \Sigma_R \tilde{\Phi} - \frac{1}{2}\bar{\Sigma}_R^c M_R^\dagger \Sigma_R + H.c. \tag{1}$$

where the super-script "c" indicates the charge conjugation. The lepton doublet $L_L = (L_{Li}) : (1,2,-1/2)$, $E_R = (E_{R_i}) : (1,1,-1)$, and Higgs doublet $\Phi = (\phi_i) : (1,2,1/2)$ ($\tilde{\Phi} = i\sigma_2\Phi^*$) have the components given by $L_{L1} = \nu_L$, $L_{L2} = e_L$, and $\phi_1 = h^+$, $\phi_2 = (v + h + iI_\phi)/\sqrt{2}$. With just one Higgs doublet, I_ϕ and h^+ are the would-be Nambu-Goldstone bosons h_z and h_w^+ "eaten" by Z and W bosons, respectively. We have

$$\begin{aligned}
&\bar{L}_L \Sigma_R \tilde{\Phi} = \bar{L}_{Li}\Sigma_{Rij}\tilde{\Phi}_{j'}\epsilon^{jj'} = -\Big(i\frac{1}{2}\bar{\nu}_L\Sigma_R^0 + \frac{1}{\sqrt{2}}\bar{e}_L\Sigma_R^-\Big)(v + h - iI_\phi) - \Big(\bar{\nu}_L\Sigma_R^+ + i\frac{1}{\sqrt{2}}\bar{e}_L\Sigma_R^0\Big)h^- \;, \\
&\bar{\Sigma}_R^c\Sigma_R = \bar{\Sigma}_{Rij}^c\Sigma_{Ri'j'}\epsilon^{ii'}\epsilon^{jj'} = \bar{\Sigma}_R^{-c}\Sigma_R^+ + \bar{\Sigma}_R^{0c}\Sigma_R^0 + \bar{\Sigma}_R^{+c}\Sigma_R^- \;.
\end{aligned} \tag{2}$$

In the above, repeated indices are summed over from 1 to 2. $\epsilon_{12} = 1$, $\epsilon_{21} = -1$ and $\epsilon_{11} = \epsilon_{22} = 0$. The neutrino and charged lepton mass matrices M_ν and M_E, in the basis $(\nu_L^c, \Sigma_R^0)^T$ and $(e_R, \Sigma_R^-)^T$, are given by

$$M_\nu = \begin{pmatrix} 0 & M_{\nu\Sigma} \\ M^T_{\nu\Sigma} & M^\dagger_R \end{pmatrix} , \quad M_E = \begin{pmatrix} M_e & M_{e\Sigma} \\ 0 & M^\dagger_R \end{pmatrix} , \tag{3}$$

where Dirac mass term $M_{\nu\Sigma} = -iY_\nu v/2$, $M_{e\Sigma} = -Y_\nu v/\sqrt{2}$ and $M_e = Y_e v/\sqrt{2}$ where v is the vacuum expectation value (VEV) of the Higgs doublet.

Note that given L_L and Σ_R representations, it is also possible to have the necessary Dirac mass term $M_{\nu\Sigma}$ from the Yukawa couplings of a quadruplet Higgs representation χ: $(1, 4, -1/2)$ of the following form,

$$L = -\bar{L}_L Y_\chi \Sigma_R \chi + H.c. \tag{4}$$

The field χ has component fields: $\chi = (\chi^+, \chi^0, \chi^-, \chi^{--})$. In tensor notation χ is a total symmetric tensor with 3 indices χ_{ijk} with i, j and k taking values 1 and 2 with

$$\chi_{111} = \chi^+ , \quad \chi_{112} = \frac{1}{\sqrt{3}}\chi^0 , \quad \chi_{122} = \frac{1}{\sqrt{3}}\chi^- , \quad \chi_{222} = \chi^{--} . \tag{5}$$

We have

$$\begin{aligned} \bar{L}_L \Sigma_R \chi &= \bar{L}_{Li} \Sigma_{Rjk} \chi_{ij'k'} \epsilon^{jj'} \epsilon^{kk'} \\ &= \bar{\nu}_L \Big(\frac{1}{\sqrt{3}} \Sigma^+_R \chi^- - i\sqrt{\frac{2}{3}} \Sigma^0_R \chi^0 + \Sigma^-_R \chi^+ \Big) + \bar{e}_L \Big(\Sigma^+_R \chi^{--} - i\sqrt{\frac{2}{3}} \Sigma^0_R \chi^- + \frac{1}{\sqrt{3}} \Sigma^-_R \chi^0 \Big) . \end{aligned} \tag{6}$$

The neutral component χ^0 can have VEV v_χ with $\chi^0 = (v_\chi + \chi_R + i\chi_I)/\sqrt{2}$. A non-zero v_χ will modify the neutrino and charged lepton mass matrices $M_{\nu\Sigma}$ and $M_{e\Sigma}$ with

$$M_{\nu\Sigma} = -i\frac{1}{2} Y_\nu v - i\frac{1}{\sqrt{3}} Y_\chi v_\chi , \quad M_{e\Sigma} = -\frac{1}{\sqrt{2}} Y_e v + \frac{1}{\sqrt{6}} Y_\chi v_\chi . \tag{7}$$

To the leading tree level light neutrino mass matrix m_ν, defined by $L_m = -\frac{1}{2}\bar{\nu}^c_L m_\nu \nu_L$ +H.c., is given by

$$m_\nu = -M^*_{\nu\Sigma} M^{-1}_R M^\dagger_{\nu\Sigma} = \Big(\frac{1}{2} Y^*_\nu v + \frac{1}{\sqrt{3}} Y^*_\chi v_\chi \Big) M^{-1}_R \Big(\frac{1}{2} Y^\dagger_\nu v + \frac{1}{\sqrt{3}} Y^\dagger_\chi v_\chi \Big) . \tag{8}$$

A model with a different Higgs quadruplet $(1, 4, 3/2)$ has also been studied where neutrino masses only arises from a dimension-7 operator [2]. This model is very different from the model we are discussing here.

In the basis where the charged lepton mass matrix is already diagonalized, the PMNS mixing matrix V[3, 4] in the charged current interaction is given by

$$\hat{m}_\nu = V^T m_\nu V , \tag{9}$$

where $\hat{m}_\nu = diag(m_1, m_2, m_3)$ is the diagonalized light neutrino mass matrix.

The introduction of quadruplet χ in the model can have interesting consequences for neutrino masses, mixing and also for lepton flavor violating (LFV) processes, $\mu \to e\gamma$ and $\mu - e$ conversion because the VEV of χ is constrained to be small which then can lead to a large Yukawa coupling Y_χ. We also found some interesting collider signatures of the doubly charged Higgs boson in the quadruplet. In the following we will study the quadruplet model in more details.

II. THE ELECTROWEAK CONSTRAINT

We have seen that in Type III seesaw, it is possible to introduce a quadruplet Higgs which give additional seesaw contributions to neutrino masses at the tree level. It is, however, well known that electroweak precision data constrain the VEV of a Higgs representation because a non-zero VEV of some Higgs may break the $SU(2)$ custodial symmetry in the SM leading to a large deviation of the ρ parameter from unity. With the constraints satisfied, the Higgs doublet and quadruplet may contribute to the neutrino mass matrix differently.

The non-zero VEV of the Higgs representation with isospin I and hypercharge Y will modify the ρ parameter at tree level with[5],

$$\rho = \frac{\sum_\alpha (I_\alpha(I_\alpha+1) - Y_\alpha^2) v_\alpha^2}{\sum_\alpha 2Y_\alpha^2 v_\alpha^2} \;. \tag{10}$$

The SM doublet Higgs alone does not lead to a deviation of ρ from unity, but the addition of a quadruplet does. For our case of one doublet and one quadruplet, we have

$$\rho = \frac{v^2 + 7v_\chi^2}{v^2 + v_\chi^2} = 1 + \frac{6v_\chi^2}{v^2+v_\chi^2} \;. \tag{11}$$

We therefore have, $\Delta\rho = 6v_\chi^2/(v^2+v_\chi^2) = 6\sqrt{2}G_F v_\chi^2$. Using experimental data $\Delta\rho = 0.0004^{+0.0029}_{-0.0011}$(95% c.l.)[6], we see that v_χ is constrained to be less than 5.8 GeV which is about 40 times smaller than that of the doublet Higgs VEV. This vast difference in Higgs VEV's indeed indicate that the Higgs doublet and quadruplet contribute to the neutrino mass matrix differently in the sense that if the Yukawa couplings Y_ν and Y_χ are the same order of magnitude, they contribute to the neutrino masses can be different by orders of magnitude. Turning this around, if both Higgs contribute to the neutrino masses with the same orders of magnitude, then the Yukawa coupling for quadruplet Y_χ can be several orders of magnitude larger than that for the doublet Y_ν.

If the seesaw mass is only from the coupling to Φ, just like Type III seesaw with one doublet, the canonical Yukawa coupling is of order $\sqrt{M_R m_\nu/v^2}$. With a M_R of order 1 TeV, the Yukawa couplings would be less than 10^{-5} with m_ν around 0.1 eV. This makes it clear that even the heavy degrees of freedom are kinematically accessible at the LHC[7], the small Yukawa couplings is hard to study their properties and their effects on LFV processes[8–10]. Although it has been shown that there are solutions with large Yukawa coupling in Type III seesaw with just one Higgs doublet[9, 11], it is interesting to see if large Yukawa couplings can more naturally manifest itself. The quadruplet with a small VEV provides such a possibility. The natural size of the Yukawa coupling Y_χ is of order $\sqrt{M_R m_\nu/v_\chi^2}$. With v_χ of order 1 GeV, Y_χ would be enhanced by about 250 times compared with Y_ν. With a smaller v_χ, Y_χ can be even larger since $Y_\chi \sim 10^{-3}(1\mathrm{GeV}/v_\chi)\sqrt{(M_R/\mathrm{TeV})(m_\nu/0.1\mathrm{eV})}$. The large Yukawa coupling Y_χ can lead to interesting phenomenology, such as the possibility of having large effects in lepton flavor violating (LFV) processes $\mu \to e\gamma$ and $\mu - e$ conversion.

III. LOOP INDUCED NEUTRINO MASS WITH JUST ONE TRIPLET LEPTON

In the Type III seesaw with just doublet Higgs, if there is just one leptonic triplet Σ_R, the resulting neutrino mass matrix m_ν for the three light neutrinos is only a rank one matrix. This implies that only one light neutrino mass is non-zero. Neutrino oscillation data show the existence of two distinct mass squared splittings, so a model with just one generation of triplet Σ_R is in conflict with data. More than one generation of Σ_R is required to have a higher ranked mass matrix to fit data. We point out that with the introduction of quadruplet χ, it is possible to raise the rank of neutrino mass matrix by including one loop contributions to the mass matrix. The tree and loop generated mass matrices together can be consistent with present data on neutrino mass and mixing. With both Higgs doublet and quadruplet, the tree level light neutrino mass matrix m_ν given in eq.(8) is still rank one if there is only one generation of Σ_R. In the following we show that the inclusion of one loop contribution can raise the rank of the mass matrix to two.

The one loop contributions involve exchange of internal quadruplet Higgs bosons and heavy leptons. In order to show this mechanism explicitly, we first identify physical Higgs states and mixing necessary for one loop generation of neutrino mass from the Higgs potential. The most general renormalizable Higgs potential is given by

$$\begin{aligned}\mathcal{V} = &-\mu^2\left(\Phi^\dagger\Phi\right) + \lambda\left(\Phi^\dagger\Phi\right)^2 + M^2\chi^\dagger\chi + \lambda^\alpha_\chi(\chi^\dagger\chi\chi^\dagger\chi)_\alpha + \lambda^\alpha_{\Phi\chi}(\Phi^\dagger\Phi\chi^\dagger\chi)_\alpha \\ &+ \left[\frac{\lambda_5}{2}(\Phi\chi)^2 + \lambda_{3\Phi}\Phi^\dagger\Phi\Phi\chi + \text{H.c}\right], \end{aligned} \tag{12}$$

where α denotes an index for $SU(2)$ contractions. The contraction of $SU(2)$ indices for each of the terms are given by

$$\begin{aligned}\chi^\dagger\chi &= \chi^*_{ijk}\chi_{ijk}, \\ (\chi^\dagger\chi\chi^\dagger\chi)_1 &= \chi^*_{ijk}\chi_{ijk}\chi^*_{lmn}\chi_{lmn}, \\ (\chi^\dagger\chi\chi^\dagger\chi)_2 &= \chi^*_{ijk}\chi_{ijn}\chi^*_{lmn}\chi_{lmk}, \\ (\chi^\dagger\chi\chi^\dagger\chi)_3 &= \chi^*_{ijk}\chi_{rjk}\chi^*_{lmn}\chi_{smn}\epsilon_{il}\epsilon_{rs}, \\ (\chi^\dagger\chi\chi^\dagger\chi)_4 &= \chi^*_{ijk}\chi_{rsk}\chi^*_{lmn}\chi_{tun}\epsilon_{il}\epsilon_{jm}\epsilon_{rt}\epsilon_{su}, \\ (\Phi^\dagger\Phi\chi^\dagger\chi)_1 &= \Phi^*_i\Phi_i\chi^*_{jkl}\chi_{jkl}, \\ (\Phi^\dagger\Phi\chi^\dagger\chi)_2 &= \Phi^*_i\Phi_j\chi^*_{jkl}\chi_{ikl}, \\ (\Phi^\dagger\Phi\chi^\dagger\chi)_3 &= \Phi^*_i\Phi_j\chi^*_{klm}\chi_{nlm}\epsilon_{ik}\epsilon_{jn}, \\ (\Phi\chi)^2 &= \Phi_i\Phi_j\chi_{i'kl}\chi_{j'mn}\epsilon_{ii'}\epsilon_{jj'}\epsilon_{km}\epsilon_{ln}, \\ \Phi^\dagger\Phi\Phi\chi &= \Phi^*_i\Phi_j\Phi_k\chi_{ij'k'}\epsilon_{jj'}\epsilon_{kk'}\ . \end{aligned} \tag{13}$$

In the above only two terms are independent for $(\chi^\dagger\chi\chi^\dagger\chi)_\alpha$. Also only two terms are independent for $(\Phi^\dagger\Phi\chi^\dagger\chi)_\alpha$. One can just take α to be equal to 1 and 2 as the independent terms for these two types of terms. In the following, we set $\lambda^3_\chi = \lambda^4_\chi = \lambda^3_{\Phi\chi} = 0$ without loss of generality.

The two terms $(\Phi\chi)^2$ and $\Phi^\dagger\Phi\Phi\chi$, break the global lepton number symmetry after the doublet and quadruplet develop non-zero VEV's. $\Phi^\dagger\Phi\Phi\chi$ then mixes Φ and χ fields. At one loop level Majorana masses will be generated for light neutrinos. There are three types of mixing terms which can be characterized to be proportional to v^2, vv_χ or v_χ^2. We have seen earlier that v is much larger than v_χ from electroweak precision data, therefore one can just keep terms proportional to v^2 for the loop generation of neutrino masses. These terms are

$$\begin{aligned} L = &-\frac{1}{2}\lambda_5 v^2\Big(\frac{1}{\sqrt{3}}\chi^+\chi^- - \frac{1}{6}(\chi_R + i\chi_I)^2\Big) \\ &- v^2\lambda_{3\Phi}\Big[\Big(\frac{1}{2}h^-\chi^+ - \frac{1}{\sqrt{3}}h^+\chi^-\Big) + \frac{1}{4\sqrt{3}}(3h + iI_\phi)(\chi_R + i\chi_I)\Big] + H.c. \end{aligned} \tag{14}$$

The above terms will generate a neutrino mass matrix proportional to $Y^*_\chi Y^\dagger_\chi$ for the first term and, $Y^*_\nu Y^\dagger_\chi$ for the second term. To have a consistent model, the elements in Y_ν are required to be much smaller than those in Y_χ. We can neglect the contribution from terms proportional to $\lambda_{3\Phi}$ in the above. Without terms proportional to $\lambda_{3\Phi}$ and v_χ, masses of component fields in χ are given by

$$\begin{aligned} m^2_{\chi_R} &\simeq M^2 + \Big(\frac{1}{2}\lambda^1_{\Phi\chi} + \frac{1}{6}\lambda^2_{\Phi\chi} - \frac{1}{3}\lambda_5\Big)v^2, \\ m^2_{\chi_I} &\simeq M^2 + \Big(\frac{1}{2}\lambda^1_{\Phi\chi} + \frac{1}{6}\lambda^2_{\Phi\chi} + \frac{1}{3}\lambda_5\Big)v^2, \\ m^2_{\chi^{\pm\pm}} &\simeq M^2 + \frac{1}{2}\left(\lambda^1_{\Phi\chi} + \lambda^2_{\Phi\chi}\right)v^2. \end{aligned} \tag{15}$$

We note that a parameter λ_5 characterizes a mass squared splitting between χ_R and χ_I, i.e., $(m^2_{\chi_R} - m^2_{\chi_I}) \simeq -(2/3)\lambda_5 v^2$. The mass matrix for singly charged scalars is given by

$$\begin{pmatrix} \chi^{+*} & \chi^- \end{pmatrix} \begin{pmatrix} M^2 + \frac{1}{2}\lambda^1_{\Phi\chi}v^2 & \frac{1}{2\sqrt{3}}\lambda_5 v^2 \\ \frac{1}{2\sqrt{3}}\lambda_5 v^2 & M^2 + \frac{1}{2}[\lambda^1_{\Phi\chi} + \frac{2}{3}\lambda^2_{\Phi\chi}]v^2 \end{pmatrix} \begin{pmatrix} \chi^+ \\ \chi^{-*} \end{pmatrix} = \begin{pmatrix} \chi_1^- & \chi_2^- \end{pmatrix} \begin{pmatrix} m^2_{\chi_1^\pm} & \\ & m^2_{\chi_2^\pm} \end{pmatrix} \begin{pmatrix} \chi_1^+ \\ \chi_2^+ \end{pmatrix}, \tag{16}$$

where

$$\begin{pmatrix} \chi_1^+ \\ \chi_2^+ \end{pmatrix} = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} \chi^+ \\ \chi^{-*} \end{pmatrix} \quad \text{with} \quad \tan 2\theta = -\frac{\sqrt{3}\,\lambda_5}{\lambda^2_{\Phi\chi}}. \tag{17}$$

The one-loop contributions to the neutrino mass matrix are calculated as[12, 13]

$$M_\nu^{\rm loop} \simeq Y_\chi^* Y_\chi^\dagger \frac{1}{8\pi^2}\left\{\frac{1}{3}m_N\Big[I\Big(\frac{m^2_{\chi_R}}{m_N^2}\Big) - I\Big(\frac{m^2_{\chi_I}}{m_N^2}\Big)\Big] - \frac{\sin(2\theta)}{\sqrt{3}}m_E\Big[I\Big(\frac{m^2_{\chi_1^+}}{m_E^2}\Big) - I\Big(\frac{m^2_{\chi_2^+}}{m_E^2}\Big)\Big]\right\}, \tag{18}$$

where m_E and m_N are masses of neutral and charged heavy leptons, and $I(x) = x\ln x/(1-x)$. The explicit dependence on λ_5 is given

$$\begin{aligned} M_\nu^{\rm loop} &\simeq \frac{\kappa}{m_N} Y_\chi^* Y_\chi^\dagger v^2, \\ \kappa &= \frac{\lambda_5}{12\pi^2}\Big\{-\frac{1}{3}\frac{m_N^2}{m^2_{\chi_R} - m^2_{\chi_I}}\Big[I\Big(\frac{m^2_{\chi_R}}{m_N^2}\Big) - I\Big(\frac{m^2_{\chi_I}}{m_N^2}\Big)\Big] + \frac{1}{2}\frac{m_N m_E}{m^2_{\chi_1^+} - m^2_{\chi_2^+}}\Big[I\Big(\frac{m^2_{\chi_1^+}}{m_E^2}\Big) - I\Big(\frac{m^2_{\chi_2^+}}{m_E^2}\Big)\Big]\Big\}. \end{aligned} \tag{19}$$

Neglecting mass splitting in a multiplet, i.e., $m_{\chi_i} = m_\chi$, $m_N = m_E$, κ is given by

$$\kappa = \frac{\lambda_5}{72\pi^2} J(m_\chi^2/m_N^2), \quad J(x) = \frac{1}{1-x} + \frac{\ln x}{(1-x)^2}, \tag{20}$$

where $J(1) = -1/2$.

Collecting contributions from the tree and loop contribution, one can write the neutrino mass matrix as

$$M_\nu^{ij} = \Big(M_\nu^{\rm tree} + M_\nu^{\rm loop}\Big)^{ij} = \frac{1}{m_N}\Big(\frac{1}{2}Y_\nu^i v + \frac{1}{\sqrt{3}}Y_\chi^i v_\chi\Big)^*\Big(\frac{1}{2}Y_\nu^j v + \frac{1}{\sqrt{3}}Y_\chi^j v_\chi\Big)^* + \frac{\kappa}{m_N}Y_\chi^{i*}Y_\chi^{j*}v^2. \tag{21}$$

The mass matrix is now rank 2 in general. This mechanism can also work even when we introduce an additional scalar doublet[14]. However such a scalar is indistinguishable from a SM Higgs doublet without additional quantum charges. The extra doublet fields can interact with other SM fermions and will induce large tree level flavor changing neutral current (FCNC) for the charged leptons. In this model, the tree level FCNC are much suppressed for charged leptons.

IV. SOME PHENOMENOLOGICAL IMPLICATIONS

A. Neutrino masses and mixing

The mass matrix obtained in the previous section, being rank two, has two non-zero eigenvalues. One of the neutrino masses is predicted to be zero. The zero mass neutrino can be m_{ν_1} or m_{ν_3} depending on whether the neutrino masses have normal or inverted hierarchy. In this section, we show that the mass matrix obtained can be made consistent with experimental data on mixing parameters.

Mass squared differences of neutrino masses and neutrino mixing have been measured to good precision[15–20]. The mass parameters are determined by global fit as[21] $\Delta m^2_{21} = (7.58^{+0.22}_{-0.26}) \times 10^{-5}$ eV^2, $|\Delta m^2_{31}| = (2.39^{+0.12}_{-0.09}) \times 10^{-3}$ eV^2; $(2.31^{+0.12}_{-0.09}) \times 10^{-3}$ eV^2) for normal (inverted) mass hiearchy. Here $\Delta m^2_{ij} = m^2_i - m^2_j$, and $m_i (i = 1\text{–}3)$. For our case, with normal hierarchy, $m^2_1 = 0$, $m^2_2 = \Delta m^2_{12}$ and $m^2_3 = \Delta m^2_{31}$. For inverted hierarchy, we then have $m^2_3 = 0$, $m^2_1 = -\Delta m^2_{31}$, and $m^2_2 = \Delta m^2_{21} - \Delta m^2_{31}$. The neutrino mixing are given by[21] $\sin^2(\theta_{23}) = 0.42^{+0.08}_{-0.03}$, $\sin^2(\theta_{12}) = 0.306^{+0.018}_{-0.005}$, and $\sin^2(\theta_{13}) < 0.028$. To the leading order, the mixing pattern can be approximated by the tribimaximal mixing matrix[22],

$$U_{\rm TB} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix}. \tag{22}$$

The light neutrino mass matrix obtained in eq.(21) can be easily made to fit data. We consider the case for $v \gg v_\chi$, such that terms proportional to v_χ can all be neglected for illustration. With this approximation, the cross term proportional to $Y^*_\nu Y^\dagger_\chi + Y^*_\chi Y^\dagger_\nu$ can be neglected.

For normal hierarchy case, by imposing the condition of the tribimaximal mixing, the Yukawa couplings can be taken to be the forms $Y_\nu \sim y_\nu (0,\ 1/\sqrt{2},\ -1/\sqrt{2})^T$, and $Y_\chi \sim y_\chi (1/\sqrt{3},\ 1/\sqrt{3},\ 1/\sqrt{3})^T$. In this case $m_3 = y^2_\nu v^2/4m_N$ and $m_2 = \kappa y^2_\chi v^2/m_N$. If the heavy neutrino mass is of order 1 TeV, $y_\nu \sim 1.80 \times 10^{-6} (m_N/1\text{TeV})^{1/2}$ and $\sqrt{\kappa} y_\chi = 0.38 \times 10^{-6} (m_N/1\text{TeV})^{1/2}$. We note that relative size of tree level and loop level contributions can be tuned by the parameter κ, which is proportional to the Higgs potential parameter λ_5. If λ_5 is small, quaruplet Yukawa coupling y_χ can be order of one. This kind of possibility is also studied in the neutrinophilic two Higgs doublet model[23]. The role of the Y_ν and Y_χ can be switched.

Similarly the model can be made consistent with inverted hierarchy. For example with $Y_\nu \sim y_\nu (\sqrt{2/3},\ -1/\sqrt{6},\ -1/\sqrt{6})^T$ and $Y_\chi \sim y_\chi (1/\sqrt{3},\ 1/\sqrt{3},\ 1/\sqrt{3})^T$, the tribimaximal mixing pattern can be realized. In this case $m_1 = y^2_\nu v^2/4m_N$ and $m_2 = \kappa y^2_\chi v^2/m_N$. If the heavy neutrino mass is of order 1 TeV, $y_\nu \sim 1.78 \times 10^{-6} (m_N/1\text{TeV})^{1/2}$ and $\sqrt{\kappa} y_\chi = 0.90 \times 10^{-6} (m_N/1\text{TeV})^{1/2}$. Again the roles of Y_ν and Y_χ can be switched.

Making perturbation to the above forms, one can get non-zero θ_{13} solutions, which is indicated by recent results at T2K[20]. For instance, for normal mass hierarchy case modifying Y_ν to be $Y^{'}_\nu = Y_\nu + \Delta Y_\nu = Y_\nu + y_\nu (a, b, c)^T$ and keep the same Y_χ, we can produce non-zero θ_{13} solutions. Using the $\Delta Y_\nu = y_\nu (-0.14, 0, 0)^T, y^2_\nu v^2/4m_N = 5.23 \times 10^{-2}$ eV, $\kappa y^2_\chi v^2/m_N = 9.14 \times 10^{-3}$eV, we obtain $m_2 = 8.78 \times 10^{-3}$, $m_3 = 4.82 \times 10^{-2}$, $\sin^2\theta_{12} = 0.323$, $\sin^2\theta_{23} = 0.44$ and $\sin^2\theta_{13} = 0.025$ which are within one σ error of the data.

For inverted mass hierarchy case, with $\Delta Y_\nu = y_\nu (-0.0095, -0.1, 0.1085)^T, y^2_\nu v^2/4m_N = 4.81 \times 10^{-2}$eV, $\kappa y^2_\chi v^2/m_N = 4.88 \times 10^{-2}$eV, we obtain $m_1 = 4.80 \times 10^{-2}$, $m_2 = 4.88 \times 10^{-2}$, $\sin^2\theta_{12} = 0.306$, $\sin^2\theta_{23} = 0.41$ and $\sin^2\theta_{13} = 0.014$ which are, again, within one σ error of the data.

Higher order loop corrections can further raise the rank of neutrino mass matrix in general. Therefore, all three light neutrinos can have non-zero masses in this model. It has been shown in Ref.[24] that the rank of the neutrino mass matrix can be rank two at two loop level even with just one triplet lepton and one Higgs doublet. However, in this case the heavy triplet lepton mass needs to be 10^{16} GeV, and hence its phenomenological consequence for collider physics is out of the scope at the LHC. Introduction of more leptonic triplet generations can also increase the rank of mass matrix too.

B. $\mu \to e\gamma$ and $\mu - e$ conversion

We now study possible effects on LFV processes $\mu \to e\gamma$ and $\mu - e$ conversion. $\mu \to e\gamma$ is induced at one loop level. There is a small contribution to $\mu - e$ conversion at the tree level due to mixing of charged light and heavy leptons. The dominant contribution come at the one loop level due to possible large Yukawa coupling Y_χ, because the size of Y_ν is constrained to be small by the absolute size of neutrino masses and the doublet Higgs VEV. The one loop induced effective Lagrangian responsible to $\mu \to e\gamma$ and $\mu - e$ conversions is given by

$$\mathcal{L} = -\overline{\psi_\mu}\sigma^{\mu\nu}(A_L P_L + A_R P_R)\psi_e F_{\mu\nu} + \sum_q eQ_q \bar{q}\gamma^\mu q \overline{\psi_\mu}\gamma_\mu P_L \psi_e B_L + \text{H.c}\ , \tag{23}$$

with Q_q being the electric charge of the q-quark, and

$$\begin{aligned}
A_L &= \frac{e}{32\pi^2} Y_\chi \Big\{ -\tfrac{1}{6}\big[\tfrac{1}{m^2_{\chi_R}} F_\Sigma(\tfrac{m_E^2}{m^2_{\chi_R}}) + \tfrac{1}{m^2_{\chi_I}} F_\Sigma(\tfrac{m_E^2}{m^2_{\chi_I}})\big] + \tfrac{2}{3}\big[\tfrac{s_\theta^2}{m^2_{\chi_1^+}} F_\chi(\tfrac{m_N^2}{m^2_{\chi_1^+}}) + \tfrac{c_\theta^2}{m^2_{\chi_2^+}} F_\chi(\tfrac{m_N^2}{m^2_{\chi_2^+}})\big] \\
&\quad + \tfrac{1}{m^2_{\chi^{++}}}\big[F_\Sigma(\tfrac{m_E^2}{m^2_{\chi^{++}}}) + 2F_\chi(\tfrac{m_E^2}{m^2_{\chi^{++}}})\big]\Big\} Y_\chi^\dagger m_\mu, \\
A_R &= \frac{m_e}{m_\mu} A_L, \\
B_L &= \frac{e}{16\pi^2} Y_\chi \Big\{ -\tfrac{1}{6}\big[\tfrac{1}{m^2_{\chi_R}} G_\Sigma(\tfrac{m_E^2}{m^2_{\chi_R}}) + \tfrac{1}{m^2_{\chi_I}} G_\Sigma(\tfrac{m_E^2}{m^2_{\chi_I}})\big] + \tfrac{2}{3}\big[\tfrac{s_\theta^2}{m^2_{\chi_1^+}} G_\chi(\tfrac{m_N^2}{m^2_{\chi_1^+}}) + \tfrac{c_\theta^2}{m^2_{\chi_2^+}} G_\chi(\tfrac{m_N^2}{m^2_{\chi_2^+}})\big] \\
&\quad + \tfrac{1}{m^2_{\chi^{++}}}\big[G_\Sigma(\tfrac{m_E^2}{m^2_{\chi^{++}}}) + 2G_\chi(\tfrac{m_E^2}{m^2_{\chi^{++}}})\big]\Big\} Y_\chi^\dagger\ ,
\end{aligned} \tag{24}$$

where

$$\begin{aligned}
F_\Sigma(z) &= \frac{z^2 - 5z - 2}{12(z-1)^3} + \frac{z\ln z}{2(z-1)^4}\ , \quad F_\chi(z) = \frac{2z^2 + 5z - 1}{12(z-1)^3} - \frac{z^2 \ln z}{2(z-1)^4}\ , \\
G_\Sigma(z) &= \frac{7z^3 - 36z^2 + 45z - 16 + 6(3z-2)\ln z}{36(1-z)^4}\ , \quad G_\chi(z) = \frac{11z^3 - 18z^2 + 9z - 2 - 6z^3 \ln z}{36(1-z)^4}\ .
\end{aligned} \tag{25}$$

The LFV $\mu \to e\gamma$ decay branching ratio is easily evaluated by

$$B(\mu \to e\gamma) = \frac{48\pi^2}{G_F^2 m_\mu^2}(|A_L|^2 + |A_R|^2). \tag{26}$$

The strength of $\mu - e$ conversion is measured by the quantity, $B^A_{\mu\to e} = \Gamma^A_{conv}/\Gamma^A_{capt} = \Gamma(\mu^- + A(N,Z) \to e^- + A(N,Z))/\Gamma(\mu^- + A(N,Z) \to \nu_\mu + A(N+1, Z-1)$. Following Ref.[25], we have

$$\frac{B^A_{\mu\to e}}{B(\mu \to e\gamma)} = R^0_{\mu\to e}(A) \left| 1 + \frac{\tilde{g}^{(p)}_{LV} V^{(p)}(A)}{A_R D(A)} + \frac{\tilde{g}^{(n)}_{LV} V^{(n)}(A)}{A_R D(A)} \right|^2\ , \tag{27}$$


Current experimental bounds

Future prospects

FIG. 1: The current and future experimental constraints on the quadruplet Yukawa couplings from $\mu \to e\gamma$ and $\mu - e$ conversion. The mass of quadruplet scalar is taken as $m_\chi = 1$ TeV.

where

$$R^0_{\mu\to e}(A) = \frac{G_F^2 m_\mu^5}{192\pi^2 \Gamma^A_{capt}} |D(A)|^2 \ . \tag{28}$$

and $\tilde{g}^{(p)}_{LV} = 2g_{LV(u)} + g_{LV(d)}, \tilde{g}^{(n)}_{LV} = g_{LV(u)} + 2g_{LV(d)}$ with $g_{LV(q)} = -eQ_q B_L/(\sqrt{2}G_F)$.

For many years, the best 90% c.l. experimental upper limit for $B(\mu \to e\gamma)$ was 1.2×10^{-11}[26]. Recently, MEG collaboration has obtained better result with the 90% c.l. upper limit[27] 2.4×10^{-12}. This new bound, as will be seen, provides important constraint for the quadruplet model discussed here. There are several measurements of $\mu - e$ conversion on various nuclei. The best bound is for Au nuclei with the 90% c.l. experimental bound given by $B^{\rm Au}_{\mu\to e} < 7 \times 10^{-13}$[28]. For Au, the relevant parameters determined by method I in Ref.[25] are given by: $D(\rm Au) = 0.189$, $V^{(p)}(\rm Au) = 0.0974, V^{(n)}(\rm Au) = 0.146$ and $R^0_{\mu\to e}(\rm Au) = 0.0036$[25]. We will use these values to study implication for our quadruplet model.

The numerical results are shown in FIG.1. In obtaining results in FIG.1, we have choosen the mass of quadruplet component field χ_i to be degenerate with a common mass of 1 TeV, and the quadruplet Yukawa coupling constant is taken as $Y_\chi \sim y_\chi(1/\sqrt{3},\ 1/\sqrt{3},\ 1/\sqrt{3})^T$ which satisfying neutrino mixing data from our previous studies for illustration. In the left panel of FIG.1, we show current experimental bounds on the quadruplet Yukawa coupling from non-observation of $\mu \to e\gamma$ and $\mu - e$ conversion as a function of ratio of triplet fermion and quadruplet scalar squared masses. We found that current constraints on quadruplet Yukawa coupling constant from $\mu - e$ conversions are weaker than that from $\mu \to e\gamma$. This is very different than the situation in a model with fourth generation where non-zero Z-penguin contribution dominates and $\mu - e$ conversion gives stronger constraints[29]. In the quadruplet model discussed here because the triplet heavy lepton Σ_R does not have hypercharge, no Z-penguin contribution and therefore $\mu - e$ conversion gives weaker constraint compared with $\mu \to e\gamma$. From the figure, we see that the quadruplet Yukawa couplings are constrained by the new MEG data to be less than 0.1 for a wide range of parameter space. As we showed y_ν is typically 10^{-6} for 1 TeV quadruplet scalars in both normal and inverted neutrino mass spectrum. The contribution from Y_ν is negligibly small. On the other hand, y_χ can be enhanced by a factor of $1/\sqrt{\kappa} \simeq 12\pi/\sqrt{-\lambda_5}$ with $m_N \sim m_\chi \sim 1$TeV. To obtain

$y_\chi \simeq 0.1$, $\lambda_5 \simeq 10^{-8}$ is required. Such a tiny λ_5 can be naturally understood as a remnant of the lepton number symmetry. The quadruplet model can have Yukawa coupling producing $\mu \to e\gamma$ closing to the present upper bound. Improved experimental limits can further constrain the model parameters.

In the right panel of FIG.1, we also show the future prospects of LFV bounds. For $\mu \to e\gamma$ we take $B(\mu \to e\gamma) = 1 \times 10^{-13}$[30] as the near future improved MEG experimental sensitivity. For $\mu - e$ conversion, there are several planed new experiments, such as Mu2E[31]/COMET[32] and PRISM[33] for $\mu - e$ conversion using Al and Ti. The sensitivities are expected to reach 10^{-16}[32] and 10^{-18}[33], respectively. For Ti and Al nuclei, the relevant parameters for our calculations are given by $D(\mathrm{Ti}) = 0.0864$, $V^{(p)}(\mathrm{Ti}) = 0.0396, V^{(n)}(\mathrm{Ti}) = 0.0468$ and $R^0_{\mu\to e}(\mathrm{Ti}) = 0.0041$, and $D(\mathrm{Al}) = 0.0362$, $V^{(p)}(\mathrm{Al}) = 0.0161, V^{(n)}(\mathrm{Al}) = 0.0173$ and $R^0_{\mu\to e}(\mathrm{Al}) = 0.0026$[25]. We see that improved $\mu \to e\gamma$ and $\mu - e$ conversion experiments can further constrain the quadruplet Yukawa coupling constant. Also note that searches for $\mu - e$ conversions can provide better constraints than that for $\mu \to e\gamma$.

C. Collider signatures of doubly charged Higgs bosons in quadruplet

Finally, we would like to make some comments about collider aspects of this model. One of the interesting feature of Type III seesaw is that the heavy leptons with a mass of a TeV or lower can be produced at the LHC. The collider phenomenology related to Type III seesaw for the heavy leptons has been studied in great detail[7, 34]. The introduction of quadruplet also leads to new phenomena in collider physics.

An interesting feature is the existence of the doubly charged particle χ^{++} in the model. Doubly charged scalar bosons also appear in other models for neutrino masses, for example, Higgs triplet in Type II seesaw Model[35, 36], and Zee-Babu model[37]. The doubly charged scalar bosons can be produced at a hadron collider through the Drell-Yan production mechanism $q\bar{q} \to \gamma, (Z^*) \to \chi^{++}\chi^{--}$[38, 39]. The vector boson fusion mechanism can also be useful to produce doubly charged particle[40] if the VEV's of the Higgs triplet v_Δ and the quadruplet v_χ are not very small. The recently results from LHC exclude doubly charged Higgs mass to be around 150 GeV if it decay predominantly through leptonic decay[41]. Unlike the Type II seesaw and Zee-Babu models, the quadruplet scalars do not have direct interaction with a pair of SM fermion and therefore cannot decay into them. The lower limit on the mass of doubly charged Higgs boson does not apply for our model.

In both Type II seesaw and the quadruplet models, if the VEV's v_Δ and v_χ are not very small, the doubly charged scalar will mainly decay into a pair of $W^\pm W^\pm$[42]. Zee-Babu model does not have such decay modes. In the case of Type II seesaw model, if $v_\Delta < 10^{-4}$ GeV, the leptonic pair decay modes will become the dominate one for the doubly charged scalar because the decay to gauge boson pair is suppressed by v_Δ while leptonic Yukawa coupling is scaled as m_ν/v_Δ. This is, however, not the case for quadruplet model.

The χ^{++} can couple to $e^+\Sigma^+$ through Yukawa coupling. Since the heavy charged lepton Σ^+ can mixing with e^+ because mixing in eq. (6) leading to $\chi^{++} \to e^+e^+$. However, the mixing in this case is proportional to $Y_\nu^2 v^2/2m_\Sigma^2$ which is small. There is another possible decay for χ^{++}. Electromagnetic loop correction[43] will make Σ^+ to be heavier than Σ^0 allowing $\Sigma^+ \to \pi^+\Sigma^0$. Then Σ^0 mixes with light neutrinos to allow $\Sigma^0 \to e^\pm W^\mp$ decay. Since the mixing between light neutrino and Σ^0 is only suppressed by a factor $Y_\nu v/m_\Sigma$, the decay mode, $\chi^{++} \to e^+\pi^+e^\pm W^\mp$ would be more important than $\chi^{++} \to e^+e^+$. This is different than Type II seesaw model in the case even the VEV's are very small.

V. CONCLUSIONS

In Type III seesaw the heavy neutrinos are contained in leptonic triplet representations. Being a triplet of $SU(2)_L$ gauge group, the heavy leptons have non-trivial structure. Concerning Yukawa interaction for seesaw mechanism, we find a new possibility of having new type of Yukawa couplings by introducing a quadruplet χ with hypercharge equal to half. When the neutral component field of χ develops a non-zero VEV, a Dirac mass terms connecting the light and heavy neutrinos can result to facilitate the seesaw mechanism. It is interesting to note that the VEV of the quadruplet Higgs is constrained to be very small from electroweak precision data. Therefore the Yukawa couplings of a quadruplet can be much larger than those in a Type III model with a Higgs doublet only. We also find that unlike the usual Type III seesaw model where at least two copies of leptonic triplets are needed, with both doublet and quadruplet Higgs representations, just one leptonic triplet is possible to have a phenomenologically acceptable model because light neutrino masses can receive sizable contributions from both the tree and one loop levels. Large Yukawa coupling may have observable effects on lepton flavor violating processes, such as $\mu \to e\gamma$ and $\mu - e$ conversion. There are also some interesting collider signatures for the doubly charged particle in the quadruplet model.

Acknowledgment:
We would like to thank Dr. Sugiyama for useful discussions. This work was partially supported by #20540282 and #21104004, NSC and NCTS of ROC, and SJTU 985 grant of China.

[1] R. Foot, H. Lew, X. G. He, G. C. Joshi, Z. Phys. **C 44**, 441 (1989).
[2] K. S. Babu, S. Nandi, Z. Tavartkiladze, Phys. Rev. **D80**, 071702 (2009).
[3] B. Pontecorvo, Sov. Phys. JETP **6**, 429 (1957) [Zh. Eksp. Teor. Fiz. **33**, 549 (1957)].
[4] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).
[5] J. F. Gunion, H. E. Haber, G. Kane and S. Dawson, The Higgs Hunter's Guide (Frontiers in Physics series, Addison-Wesley, Reading, MA, 1990).
[6] K. Nakamura *et al.,* [Particle Data Group], J. Phys. G **37**, 075021 (2010).
[7] B. Bajc, G. Senjanovic, JHEP **0708**, 014 (2007), B. Bajc, M. Nemevsek, G. Senjanovic, Phys. Rev. **D76**, 055011 (2007), R. Franceschini, T. Hambye, A. Strumia, Phys. Rev. D **78**, 033002 (2008), T. Li and X. G. He, Phys. Rev. D **80**, 093003 (2009) [arXiv:0907.4193 [hep-ph]].
[8] A. Abada, C. Biggio, F. Bonnet, M. B. Gavela, T. Hambye, Phys. Rev. D **78**, 033007 (2008).
[9] X. -G. He, S. Oh, JHEP **0909**, 027 (2009).
[10] J. F. Kamenik, M. Nemevsek, JHEP **0911**, 023 (2009).
[11] X. -G. He, S. Oh, J. Tandean, C. -C. Wen, Phys. Rev. D **80**, 073012 (2009).
[12] E. Ma, Phys. Rev. D **73**, 077301 (2006).
[13] E. Ma and D. Suematsu, Mod. Phys. Lett. A **24**, 583 (2009).
[14] A. Ibarra, C. Simonetto, [arXiv:1107.2386 [hep-ph]].
[15] B. T. Cleveland *et al.,*, Astrophys. J. **496**, 505 (1998), W. Hampel *et al.,* [GALLEX Collaboration], Phys. Lett. B **447**, 127 (1999), B. Aharmim *et al.,* [SNO Collaboration], Phys. Rev. Lett. **101**, 111301 (2008), J. N. Abdurashitov *et al.,* [SAGE Collaboration], Phys. Rev. C **80**, 015807 (2009), K. Abe *et al.,* [Super-Kamiokande Collaboration], Phys. Rev. D **83**, 052010 (2011), G. Bellini *et al.,* [Borexino Collaboration], [arXiv:1104.1816 [hep-ex]].
[16] R. Wendell *et al.,* [Kamiokande Collaboration], Phys. Rev. D **81**, 092004 (2010).

[17] M. H. Ahn *et al.,* [K2K Collaboration], Phys. Rev. D **74**, 072003 (2006), P. Adamson *et al.,* [The MINOS Collaboration], Phys. Rev. Lett. **106**, 181801 (2011).

[18] M. Apollonio *et al.,* [CHOOZ Collaboration], Eur. Phys. J. C **27**, 331 (2003).

[19] A. Gando *et al.,* [The KamLAND Collaboration], Phys. Rev. D **83**, 052002 (2011).

[20] T2K Collaboration, Phys. Rev. Lett. **107**, 041801 (2011) [arXiv:1106.2822].

[21] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo, A. M. Rotunno, [arXiv:1106.6028 [hep-ph]].

[22] P. F. Harrison, D. H. Perkins, W. G. Scott, Phys. Lett. B **530**, 167 (2002), Z. -z. Xing, Phys. Lett. B **533**, 85-93 (2002), X. G. He, A. Zee, Phys. Lett. B **560**, 87-90 (2003).

[23] N. Haba and K. Tsumura, JHEP **1106**, 068 (2011).

[24] Y. Liao, J. -Y. Liu, G. -Z. Ning, Phys. Rev. **D79**, 073003 (2009).

[25] R. Kitano, M. Koike and Y. Okada, Phys. Rev. D **66**, 096002 (2002) [Erratum-ibid. D **76**, 059902 (2007)].

[26] M. L. Brooks *et al.,* [MEGA Collaboration], Phys. Rev. Lett. **83**, 1521 (1999) [arXiv:hep-ex/9905013].

[27] J. Adam et *al.,* MEG Collaboration, [arXiv:1107.5547].

[28] W. H. Bertl *et al.,* [SINDRUM II Collaboration], Eur. Phys. J. C **47**, 337 (2006).

[29] N. Deshpande, T. Enkhbat, T. Fukuyama, X. G. He, L. H. Tsai and K. Tsumura, [arXiv:1106.5085 [hep-ph]].

[30] Research Proposal to INFN, "The MEG experiment: search for the $\mu^+ \to e^+\gamma$ decay at PSI", September 2002.

[31] J. P. Miller [Mu2E collabaration], *Proposal to Search for* $\mu^- N \to e^- N$ *with a Single Event Sensitivity Below* 10^{-16}.

[32] Y. Kuno *et.al.,* [COMET collaboration], *An Experimental Search for lepton Flavor Violating* $\mu - e$ *Conversion at Sensitivity of* 10^{-16} *with a Slow-Extracted Bunched Beam.*

[33] Y. Kuno *et.al.,* [PRISM/PRIME Group], Letter of Intent, *An Experimental Search for a* $\mu - e$ *Conversion at Sensitivity of the Order of* 10^{-18} *with a Highly Intense Muon Source: PRISM.*

[34] A. Arhrib, B. Bajc, D. K. Ghosh, T. Han, G. -Y. Huang, I. Puljak, G. Senjanovic, Phys. Rev. **D82**, 053004 (2010).

[35] W. Konetschny and W. Kummer, Phys. Lett. B **70**, 433 (1977); T. P. Cheng and L. F. Li, Phys. Rev. D **22**, 2860 (1980); M. Magg and C. Wetterich, Phys. Lett. B **94**, 61 (1980).

[36] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980).

[37] A. Zee, Nucl. Phys. B **264**, 99 (1986). K. S. Babu, Phys. Lett. B **203**, 132 (1988).

[38] A. G. Akeroyd, M. Aoki, Phys. Rev. D **72**, 035011 (2005).

[39] A. G. Akeroyd, C. -W. Chiang, Phys. Rev. D **80**, 113010 (2009); A. G. Akeroyd, C. -W. Chiang, N. Gaur, JHEP **1011**, 005 (2010).

[40] S. Godfrey, K. Moats, Phys. Rev. D **81**, 075026 (2010).

[41] CMS-PAS-HIG-11-001. For recent results, see also slides presented by A. Kumar at EPS2011 conference; `http://indico.in2p3.fr/contributionDisplay.py?sessionId=6&contribId=817&confId=5116`.

[42] P. Fileviez Perez, T. Han, G. -y. Huang, T. Li, K. Wang, Phys. Rev. D **78**, 015018 (2008).

[43] M. Cirelli, A. Strumia, New J. Phys. **11**, 105005 (2009).